

INTERFERENCE FRAGMENTATION FUNCTIONS AND VALENCE QUARK SPIN DISTRIBUTIONS IN THE NUCLEON*

R. L. Jaffe^{a,b,†}, Xuemin Jin^{a,‡} and Jian Tang^{a,§}

^a*Center for Theoretical Physics
Laboratory for Nuclear Science
and Department of Physics
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139*

^b*RIKEN BNL Research Center
Brookhaven National Laboratory, Upton, NY 11973*

(MIT-CTP-2690 hep-ph/9710561 October 1997)

Abstract

We explore further applications of the twist-two quark interference fragmentation functions introduced earlier. We show that semi-inclusive production of two pions in the current fragmentation region in deep inelastic scattering of a longitudinally polarized electron on a longitudinally polarized nucleon can provide a probe of the valence quark spin (or helicity difference) distribution in the nucleon.

Submitted to: *Physical Review D*

[†]Email address: jaffe@mitlns.mit.edu

[‡]Email address: jin@ctpa02.mit.edu

[§]Email address: jtang@mitlns.mit.edu

*This work is supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under cooperative research agreement #DF-FC02-94ER40818, and by the RIKEN BNL Research Center.

Measurements of quark and gluon distributions within hadrons provide us with valuable information about the nonperturbative nature of the quarks and gluons inside the hadrons. In a recent Letter [1], we have studied semi-inclusive production of two pions in the current fragmentation region in deep inelastic scattering on a transversely polarized nucleon. This may provide a practical way to measure the quark transversity distribution in the nucleon, which has proved difficult to access experimentally [2–8]. In this paper, we extend our study to the case of a longitudinally polarized electron beam scattering off a longitudinally polarized nucleon target. We show that the interference between the s - and p -wave of the two-pion system around the ρ can provide an asymmetry [Eq. (6)] which is sensitive to the valence quark spin distribution in the nucleon. Note that the asymmetry would vanish by C-invariance if the two pions were in a charge conjugation eigenstate. Hence there is no effect in regions of the $\pi\pi$ mass dominated by a single resonance. Significant effects are possible, however, in the ρ mass region where the s - and p -wave production channels are both active and provide exactly the charge conjugation mixing necessary.

The asymmetry we obtain throws pions of one charge forward along the fragmentation axis relative to pions of the other charge. If one integrated over the other kinematic variables, whatever result persisted would appear as a difference between the π^+ and π^- fragmentation functions correlated with the valence quark spin distributions. This effect in single pion fragmentation was proposed and studied some years ago by Frankfurt *et al.* [9] and Close and Milner [10]. The asymmetry we describe is therefore one particular contribution to this more general effect, with the advantage that it can be characterized in terms of $\pi\pi$ phase shifts and two particle fragmentation functions that appear in other hard processes.

Consider the semi-inclusive deep inelastic scattering process: $\vec{e}\vec{N} \rightarrow e'\pi^+\pi^-X$. We define the kinematics as follows. The four-momenta of the initial and final electron are $k = (E, \vec{k})$ and $k' = (E', \vec{k}')$, and the nucleon's momentum is P_μ . The momentum of the virtual photon is $q = k - k'$, and $Q^2 = -q^2 = -4EE' \sin^2 \theta/2$, where θ is the electron scattering angle. The standard variables in DIS, $x = Q^2/2P \cdot q$ and $y = P \cdot q/P \cdot k$, are adopted. We work at low $\pi\pi$ invariant mass, where only the s - and p -waves are significant. The $\sigma[(\pi\pi)_{l=0}^{I=0}]$ and $\rho[(\pi\pi)_{l=1}^{I=1}]$ resonances are produced in the current fragmentation region with momentum P_h and momentum fraction $z = P_h \cdot q/q^2$.¹ The invariant squared mass of the two-pion system is $m^2 = (k_+ + k_-)^2$, with k_+ and k_- the four-momentum of π^+ and π^- , respectively. The decay polar angle in the rest frame of the two-pion system is denoted by Θ . Note that the azimuthal angle ϕ of the two-pion system does not figure in present analysis and can be integrated out.

Following Ref. [1], we use a collinear approximation, *i.e.*, $\theta \approx 0$ for simplicity, and work only to the leading twist (the complete analysis will be published elsewhere [11]). Invoking the helicity density matrix formalism developed in Refs. [12,8], we factor the process into various basic ingredients expressed as helicity density matrices:

¹We recognize that the $\pi\pi$ s -wave is not resonant in the vicinity of the ρ and our analysis does not depend on a resonance approximation. For simplicity we refer to the non-resonant s -wave as the “ σ ”.

$$\left[\frac{d^5\sigma}{dx dy dz dm^2 d\cos\Theta} \right]_{H'H} = \mathcal{F}_{H'H}^{h_1 h'_1} \left[\frac{d^2\sigma(eq \rightarrow e'q')}{dx dy} \right]_{h'_1 h_1}^{h_2 h'_2} \left[\frac{d^2\hat{\mathcal{M}}}{dz dm^2} \right]_{h'_2 h_2}^{H_1 H'_1} \left[\frac{d\mathcal{D}}{d\cos\Theta} \right]_{H'_1 H_1}, \quad (1)$$

where $h_i(h'_i)$ and $H(H')$ are indices labeling the helicity states of quark and nucleon, and $H_1(H'_1)$ labeling the helicity state of the resonance (σ, ρ) . Physically, the four factors on the right-hand side of Eq. (1) represent the $N \rightarrow q$ distribution function, the hard partonic $eq \rightarrow e'q'$ cross section, the $q \rightarrow (\sigma, \rho)$ fragmentation, and the decay $(\sigma, \rho) \rightarrow \pi\pi$, respectively. In order to incorporate the final state interaction, we have separated the $q \rightarrow \pi^+\pi^-$ fragmentation process into two steps. First, the quark fragments into the resonance (σ, ρ) , then the resonance decays into two pions (see Figure 1 of Ref. [1]).

The s - p interference fragmentation function describes the emission of a $\rho(\sigma)$ with helicity H_1 from a quark of helicity h_2 , followed by absorption of $\sigma(\rho)$, with helicity H'_1 forming a quark of helicity h'_2 . Imposing various symmetry restrictions, the interference fragmentation can be cast into a double density matrix [1]

$$\begin{aligned} \frac{d^2\hat{\mathcal{M}}}{dz dm^2} = & \Delta_0(m^2) \left\{ I \otimes \bar{\eta}_0 \hat{q}_I(z, m^2) + (\sigma_+ \otimes \bar{\eta}_- + \sigma_- \otimes \bar{\eta}_+) \delta \hat{q}_I(z, m^2) \right\} \Delta_1^*(m^2) \\ & + \Delta_1(m^2) \left\{ I \otimes \eta_0 \hat{q}_I(z, m^2) + (\sigma_- \otimes \eta_+ + \sigma_+ \otimes \eta_-) \delta \hat{q}_I(z, m^2) \right\} \Delta_0^*(m^2), \end{aligned} \quad (2)$$

where $\sigma_{\pm} \equiv (\sigma_1 \pm i\sigma_2)/2$ with $\{\sigma_i\}$ the usual Pauli matrices. The η 's are 4×4 matrices in (σ, ρ) helicity space with nonzero elements only in the first column, and the $\bar{\eta}$'s are the transpose matrices ($\bar{\eta}_0 = \eta_0^T, \bar{\eta}_+ = \eta_-^T, \bar{\eta}_- = \eta_+^T$), with the first rows $(0, 0, 1, 0)$, $(0, 0, 0, 1)$, and $(0, 1, 0, 0)$ for $\bar{\eta}_0$, $\bar{\eta}_+$, and $\bar{\eta}_-$, respectively. The final state interactions between the two pions are included explicitly in $\Delta_0(m^2) = -i \sin \delta_0 e^{i\delta_0}$ and $\Delta_1(m^2) = -i \sin \delta_1 e^{i\delta_1}$, with δ_0 and δ_1 the strong interaction $\pi\pi$ phase shifts. Here we have suppressed the m^2 dependence of the phase shifts. Note that $\hat{q}_I$ and $\delta \hat{q}_I$ depend on both m^2 and z . We expect that the principal m^2 dependence will enter through the final state factors Δ_0 and Δ_1 . To preserve clarity, the Q^2 dependence of the fragmentation functions has been suppressed. Henceforth we suppress the m^2 dependence as well.

The decay density matrix, quark distribution function, and hard scattering cross section have been given explicitly in Ref. [1]. For completeness we quote the results relevant to the present analysis. The interference part of the decay density matrix (after integration over the azimuthal angle ϕ) is

$$\frac{d\mathcal{D}}{d\cos\Theta} = -\frac{\sqrt{3}}{2\pi m} \cos\Theta (\bar{\eta}_0 + \eta_0). \quad (3)$$

The quark distribution function $\mathcal{F}$ is expressed as [8]

$$\mathcal{F} = \frac{1}{2}q(x) I \otimes I + \frac{1}{2}\Delta q(x) \sigma_3 \otimes \sigma_3 + \frac{1}{2}\delta q(x) (\sigma_+ \otimes \sigma_- + \sigma_- \otimes \sigma_+), \quad (4)$$

where the first matrix in the direct product is in the nucleon helicity space and the second in the quark helicity space. Here $q(x)$, $\Delta q(x)$, and $\delta q(x)$ are the spin average, spin, and transversity distribution functions, respectively, and their dependence on Q^2 has been suppressed. Finally, for a longitudinally polarized electron beam, the hard scattering cross section is given by [8]

$$\frac{d^2\sigma_{\pm}(eq \rightarrow e'q')}{dx dy} = \frac{e^4 e_q^2}{8\pi Q^2} \left[\frac{1 + (1-y)^2}{2y} (I \otimes I + \sigma_3 \otimes \sigma_3) + \frac{2(1-y)}{y} (\sigma_+ \otimes \sigma_- + \sigma_- \otimes \sigma_+) \right. \\ \left. \pm \frac{2-y}{2} (\sigma_3 \otimes I + I \otimes \sigma_3) \right], \quad (5)$$

in the collinear approximation, where the $\pm$ sign refers to the initial electron helicity. Here e_q is the charge fraction carried by a quark, and we have integrated out the azimuthal angle of the scattering plane.

The cross section can be obtained by putting all the ingredients together. To facilitate our discussions, we define forward and backward cross sections, $d\sigma^F$ and $d\sigma^B$, where the Θ dependence has been integrated over the forward ($0 \leq \Theta \leq \pi/2$) and backward ($\pi/2 \leq \Theta \leq \pi$) hemisphere in the two-pion rest frame, respectively. For a longitudinally polarized nucleon target with a longitudinally polarized electron beam, we then obtain the following double asymmetry

$$\mathcal{A}_{\uparrow\downarrow}^{\text{FB}} \equiv \frac{(d\sigma_{\uparrow\uparrow}^F - d\sigma_{\uparrow\uparrow}^B) - (d\sigma_{\uparrow\downarrow}^F - d\sigma_{\uparrow\downarrow}^B)}{(d\sigma_{\uparrow\uparrow}^F + d\sigma_{\uparrow\uparrow}^B) + (d\sigma_{\uparrow\downarrow}^F + d\sigma_{\uparrow\downarrow}^B)} = -\frac{\sqrt{3}y(2-y)}{1 + (1-y)^2} \sin\delta_0 \sin\delta_1 \cos(\delta_0 - \delta_1) \\ \times \frac{\sum_a e_a^2 \Delta q_a(x) \hat{q}_I^a(z)}{\sum_a e_a^2 q_a(x) [\sin^2\delta_0 \hat{q}_0^a(z) + \sin^2\delta_1 \hat{q}_1^a(z)]}, \quad (6)$$

where $\hat{q}_0$ and $\hat{q}_1$ are spin-average fragmentation functions for the σ and ρ resonances, respectively. The arrows $\uparrow\uparrow$ and $\uparrow\downarrow$ ($\uparrow$ and $\downarrow$) indicate the nucleon's (electron's) polarization along and opposite to the beam direction. Note that in Eq. (6) the electron's polarization is fixed and the asymmetry requires flipping the nucleon's polarization. This result also holds when the electron and nucleon polarizations are exchanged.

The flavor structure of the asymmetry $\mathcal{A}_{\uparrow\downarrow}^{\text{FB}}$ is quite simple. Isospin symmetry gives $\hat{u}_I = -\hat{d}_I$ and $\hat{s}_I = 0$, and charge conjugation requires $\hat{q}_I^a = -\hat{\bar{q}}_I^a$. This implies $\sum_a e_a^2 \Delta q_a \hat{q}_I^a = [4/9 \Delta u_v - 1/9 \Delta d_v] \hat{u}_I$, where $\Delta q_v \equiv \Delta q - \Delta \bar{q}$. Therefore $\mathcal{A}_{\uparrow\downarrow}^{\text{FB}}$ is sensitive to the valence quark spin distribution. Note that the interference between the two partial waves makes the interference fragmentation function $\hat{q}_I^a(z)$ charge conjugation odd and accesses the valence quark spin distribution. The role of final state interactions is quite different here than in Ref. [1]. Here the effect persists as long as two partial waves of opposite C-parity are active, whether or not they are out of phase. This is evidenced in Eq. (6) by the factor $\cos(\delta_0 - \delta_1)$. Note that if either δ_0 or δ_1 goes to zero and thus only one partial wave is active, the asymmetry vanishes as required by charge conjugation. In Fig. 1 we have plotted the factor, $\sin\delta_0 \sin\delta_1 \cos(\delta_0 - \delta_1)$, as a function of the two-pion invariant mass m . It is positive over the ρ region and hence the effect does not average to zero over this region. This differs from the case of the transversity asymmetry derived in Ref. [1], where the interference averages to zero over the ρ region due to a factor $\sin(\delta_0 - \delta_1)$. We also see from Fig. 1 that the interference peaks near the ρ mass, indicating that an optimal signal would be in the vicinity of $m \sim m_\rho$. It is unclear at this stage whether the effect would survive averaging over the z dependence of the interference fragmentation function.

A similar forward-backward asymmetry appears in the unpolarized process




FIG. 1. The factor, $\sin \delta_0 \sin \delta_1 \cos(\delta_0 - \delta_1)$, as a function of the invariant mass m of two-pion system. The data on $\pi\pi$ phase shifts are taken from Ref. [13].

$$\mathcal{A}^{\text{FB}} \equiv \frac{d\sigma^{\text{F}} - d\sigma^{\text{B}}}{d\sigma^{\text{F}} + d\sigma^{\text{B}}} = -\sqrt{3} \sin \delta_0 \sin \delta_1 \cos(\delta_0 - \delta_1) \times \frac{\sum_a e_a^2 q_a(x) \hat{q}_I^a(z)}{\sum_a e_a^2 q_a(x) [\sin^2 \delta_0 \hat{q}_0^a(z) + \sin^2 \delta_1 \hat{q}_1^a(z)]}. \quad (7)$$

Isospin symmetry and charge conjugation again dictate that $\sum_a e_a^2 q_a \hat{q}_I^a = [4/9 u_v - 1/9 d_v] \hat{u}_I$ for $\pi^+\pi^-$ system, where $q_v \equiv q - \bar{q}$. Given that the valence quark distributions u_v and d_v are known experimentally, this asymmetry can be used to determine the interference fragmentation function $\hat{q}_I^a(z)$ and hence isolate the valence quark spin distribution from $\mathcal{A}_{\uparrow\downarrow}^{\text{FB}}$. This point becomes more transparent when $\mathcal{A}_{\uparrow\downarrow}^{\text{FB}}$ is combined with $\mathcal{A}^{\text{FB}}$. In this case, one can find an asymmetry *independent* of the interference fragmentation functions:

$$\frac{(d\sigma_{\uparrow\uparrow}^{\text{F}} - d\sigma_{\uparrow\uparrow}^{\text{B}}) - (d\sigma_{\uparrow\downarrow}^{\text{F}} - d\sigma_{\uparrow\downarrow}^{\text{B}})}{(d\sigma_{\uparrow\uparrow}^{\text{F}} - d\sigma_{\uparrow\uparrow}^{\text{B}}) + (d\sigma_{\uparrow\downarrow}^{\text{F}} - d\sigma_{\uparrow\downarrow}^{\text{B}})} = \frac{y(2-y)}{1+(1-y)^2} \frac{4\Delta u_v - \Delta d_v}{4u_v - d_v}. \quad (8)$$

So, one may use $\pi^+\pi^-$ production on both nucleon and deuteron targets to measure Δu_v and Δd_v . Although the results in Eq. (8) are independent of z , one should keep the cross section differential in z to avoid possibly washing out the effect.

Refs. [9,10] explain how to use single meson (pion or kaon) production in deep inelastic scattering to measure the valence quark spin distribution in the nucleon. It is clear from Eq. (8) that our asymmetry is a particular contribution to the ones described in Refs. [9,10]. Ours is perhaps more under control since it is differential in m^2 and expressed in terms

of $\pi\pi$ phase shifts. The single particle asymmetry makes use of a larger data set. These independent ways of measuring the valence quark spin distributions should both be pursued. In addition, the asymmetries we have studied are sensitive to two particle interference fragmentation functions which may be interesting quantities in their own right. These measurements may be carried out in facilities such as HERMES at HERA and COMPASS at CERN, both of which have sensitivity to the hadronic final state in electron scattering.

To summarize, we have discussed the applications of the twist-two interference quark fragmentation functions introduced previously to the case of longitudinally polarized electron beam and longitudinally polarized nucleon target. We obtain two asymmetries: one provides a probe of the valence quark spin distribution, and the other can be used to extract the interference fragmentation functions.

We would like to thank John Collins for pointing out an error in the earlier version of this paper.

REFERENCES

- [1] R. L. Jaffe, Xuemin Jin, and Jian Tang, MIT-CTP-2672, hep-ph/9709322.
- [2] J. Ralston and D. E. Soper, Nucl. Phys. **B152**, 109 (1979).
- [3] X. Artru and M. Mekhfi, Z. Phys. **C 45**, 669 (1990).
- [4] R. L. Jaffe and X. Ji, Phys. Rev. Lett, **67**, 552 (1991).
- [5] J. C. Collins, Nucl. Phys. **B396**, 161 (1993).
- [6] X. Ji, Phys. Rev. **D 49**, 114 (1994).
- [7] J. C. Collins, S. F. Heppelmann, and G. A. Ladinsky, Nucl. Phys. **B420**, 565 (1994).
- [8] R. L. Jaffe, Phys. Rev. **D 54**, 6581 (1996).
- [9] L. L. Frankfurt *et al.*, Phys. Lett. **B230**, 141 (1989).
- [10] F. E. Close and R. G. Milner, Phys. Rev. **D 44**, 3691 (1991).
- [11] R. L. Jaffe, X. Jin, and J. Tang, in preparation.
- [12] K. Chen, G. R. Goldstein, R.L. Jaffe, X. Ji, Nucl. Phys. **B445**, 380 (1995).
- [13] P. Estabrooks and A. D. Martin, Nucl. Phys. **B79**, 301 (1974).